








- 1. To best control your pain, take ibuprofen (Motrin®, Advil®) every 6 hours.
- 2. Start taking ibuprofen (Motrin®, Advil®) immediately following surgery.
- 3. Use acetaminophen (Tylenol®) in between doses of ibuprofen (Motrin®, Advil®) if needed.
- 4. Only take **oxycodone/hydrocodone** for severe pain. We do not expect that you will need to take any **oxycodone/hydrocodone**.
- 5. Do not take more than **4000mg of Tylenol**® in a 24 hour period.







Left over Oxycodone/hydrocodone pills? Get of them! Please get unused opioids out of your home. Find local dropbox here:

3) (1/2 3) (1/2 3)

www.awarerx.pharmacy

Fact: 4 out of 5 heroin users started out misusing prescription painkillers.



SIDE EFFECTS INCLUDE:

Nausea, Vomiting, Constipation Headache, Legal Impairment & Risk of Addiction If you need more pain relief than the routine shown on above, consider taking at anytime:

Recommended to start with 1/2 pill and only take more if needed.

OXYCODONE/ HYDROCODONE 5MG (1/2-1 PILL EVERY 4-6 HRS.)

DISCLAIMER: The information provided regarding the drug regimen is for general informational purposes only. It is essential that you consult with your healthcare provider or doctor before starting, changing, or stopping any medication or drug regimen. Your primary case physician or doctor is the best person to assess your overall medical condition. Always follow the recommendations of your healthcare provider or doctor and contact them immediately if you have any concerns or questions related to your health or medications.

